Buellia ocellata, new report for New England 430–447
Bulbostylis capillaris distribution in New England 237–360, map 269
Burnett, Alison 415–429

Camilo, Rubén Martínez 187–196
Camissonia 395–414; ***C. dominguez-escalantorum***, sp. nov. 396–399; ***C. holanderi***, sp. nov. 399–402; key to related species 396
Campanula 161–186; *C. biflora* 168; *C. intermedia* 168; *C. ludoviciana* 167–168; *C. montevidensis* 168
Campanula punctata, new record for Connecticut 459–463
Canada 101–107, 125–136
Cardamine flexuosa, new record for Connecticut 459–463
Carex distribution in New England 237–360, maps 269–329
Carolina bay 137–144
Catling, Paul M., Gisèle Mitrow, and Lynn Black. Analysis of stem color and correlated morphological characters for grouping *Phragmites* (Poaceae) taxa in eastern Ontario. 125–136
Ceranthera linearifolia 387–394
Ceska, Adolf 101–107
Ceska, Oldriska 101–107
Chamaedorea 187–196; *C. glaucifolia* 189–191; *C. tuerckheimii* 191–193
Checklist for Contributors to *Rhodora* 117–124
Chiapas 187–196
China 448–455
Cladistics, *Nuphar* 1–95
Cladium mariscoides distribution in New England 237–360, map 329
Cladonia symphycarpia, new report for New England 430–447
Clairvillea 1–95, C. *lutea* 26
Clarification of the type locality of *Dicerandra linearifolia* (Labiatae). 387–394
Common reed 96–100, 125–136, 415–429
Comparison of morphological variation indicative of ploidy level in *Phragmites australis* (Poaceae) from eastern North America. 415–429
Comparison of pollen and stomata in late-glacial and early-Holocene lake sediments from eastern Massachusetts. 225–229 (New England Note)
Compositae (Asteraceae) 395–414
Connecticut 459–463
Coronilla scorpioides, new record for Connecticut 459–463
Cruciferae (Brassicaceae) 395–414
Cucumis 161–186; C. *odoratissimus* 168–169
Cyperaceae 237–360, distribution maps 268–358
Cyperus distribution in New England 237–360, maps 329–335
Cypripedium parviflorum, Holocene biogeography and genetic diversity 369–386

Dianthera 161–186; *D. laete-virens* 169
Dicerandra linearifolia (Labiatae), clarification of the type locality 387–394, distribution map 388
Dicliptera 161–186; *D. halei* 169–170; *D. viridiflora* 169
Disjunct, *Myriophyllum quitense* 101–107
Distribution of Cyperaceae in New England 237–360
Distribution of *Nuphar*, global 1–95, 13 maps 29–84
Dryas integrifolia, Holocene biogeography and genetic diversity 369–386
Dulichium arundinaceum var. *arundinaceum* distribution in New England 237–360, map 335
Dysmicodon 161–186; *D. californicum* 168; *D. ovatum* 168

Edaphic factors influence lichen species presence 430–447
Eleocharis distribution in New England 237–360, maps 335–342
Elliott, Stephen 387–394

Endangered species, Mexico 187–196
Erigeron 395–414; ***E. katiae***, sp. nov. 412–413
Eriogonum 161–186; *E. setaceum* 170–171; *E. tenellum* 170–171, var. *caulescens* 171, var. *ramosissimum* 171
Eriophorum distribution in New England 237–360, maps 342–344
Eupatorium 137–144, 145–160
Eupatorium leucolepis, ***E. paludicola***, *E. mohrii* × *E. paludicola* 137–144, key 143
Eupatorium novae-angliae, comparison with *E. leucolepis*, *E. paludicola*, *E. perfoliatum* 145–160
Eupatorium paludicola (Asteraceae): A new species from the coastal plain of North and South Carolina. 137–144, key 143

Fimbristylis autumnalis distribution in New England 237–360, map 344
Flora of: China 448–455; Connecticut 459–463; New England: Cyperaceae 237–360
Forest ecology 197–221
Forest vegetation of the Ozark Mountains in northwestern Arkansas. Upland, 197–221
Foster, David R. 225–229
Freshwater red algal species, ***Tuomeya americana***, comb. nov. 464–467
Fuirena pumila, *F. squarrosa* distribution in New England 237–360, maps 344
Fuscopannaria praetermissa, new report for Maine 430–447

Gatesia 161–186; *G. laete-virens* 169
Genetic diversity 369–386
Georgia pine barrens 387–394
Geum peckii, Holocene biogeography and genetic diversity 369–386
Glacial (late) lake sediments from eastern Massachusetts, pollen and stomata in 225–229
Glennon, Kelsey 415–429
Glyceria maxima, new record for Connecticut 459–463
Guard cell size, *Phragmites* 415–429

Haloragaceae 101–107
Hansen, Barbara C. S. 225–229
Harris, Tanner B., Fred C. Olday, and Nishanta Rajakaruna. Lichens of Pine Hill, a peridotite outcrop in eastern North America. 430–447
Holocene 225–229, 369–386
Holocene lake sediments from eastern Massachusetts, pollen and stomata in 225–229
Huck, Robin B. Clarification of the type locality of *Dicerandra linearifolia* (Labiatae). 387–394
Huebner, Cynthia D. 197–221
Hunter, Kimberly L. 415–429
Hunter, Richard B. 415–429
Hymenoxys 395–414; ***H. argentea*** var. ***thoreauensis***, var. nov. 410–411

Iglesias, Carlos 187–196
Introduced species: *Phragmites* 96–100, 125–136, 415–429
Invasive plant identification 125–136
Invasive species, *Phragmites* 96–100, 125–136, 415–429
ITS sequences 145–160

John L. Riddell, North American taxa published in 1853 161–186
Justicia 161–186; *J. laetevirens* 169; *J. viridiflora* 169

Kyllingia gracillima distribution in New England 237–360, map 345

Labiatae 387–394
Larson, Joseph S. and Karen B. Searcy. *Lemna minuta* (Lemnaceae) discovered in Massachusetts. 456–458 (New England Note)
Lathyrus japonicus, Holocene biogeography and genetic diversity 369–386

LeBlond, Richard J. 137–144, 145–160
LeBlond, Richard J., Edward E. Schilling, Richard D. Porcher, Bruce A. Sorrie, J. F. Townsend, Patrick D. McMillan, and Alan S. Weakley. ***Eupatorium paludicola*** (Asteraceae): A new species from the coastal plain of North and South Carolina. 137–144
Legousia 161–186; *L. biflora* 168
Lemanea [*'Lemania'*] *americana* 464–467
Lemna minuta (Lemnaceae) discovered in Massachusetts. 456–458 (New England Note)
Lepidium 395–414; ***L. montanum*** var. ***todiltoensis***, var. nov. 407–409
Lichen ecology 430–447
Lichens of Pine Hill, a peridotite outcrop in eastern North America. 430–447
Lipocarpha micrantha distribution in New England 237–360, map 345
Lobaria, new report for ultramafic soils worldwide 430–447
Lysimachia 161–186; *L. angustifolia* 171; *L. carpenteri* 171–172; *L. ciliata* var. *angustifolia* 172, var. *heterophylla* 172; *L. graminea* 172; *L. heterophylla* 171; *L. lanceolata* 171–172, var. *angustifolia* 172, var. *heterophylla* 172; *L. procumbens* 172–173; *L. radicans* 173

Maine, lichens 430–447
Massachusetts 96–100, 225–229, 456–458
McAlpine, Donald F., Gart Bishop, Oldriska Ceska, Michael L. Moody, and Adolf Ceska. Andean Watermilfoil, *Myriophyllum quitense* (Haloragaceae), in the Saint John River estuary system, New Brunswick, Canada. 101–107 (Note)
McMillan, Patrick D. 137–144
Meléndez, Nayely Martínez 187–196
Melothria 161–186; *M. pendula* 173; *M. punctata* 173–174; *M. repanda* 173
Mexico, palms 187–196
Migration of plants 369–386
Mitrow, Gisèle 125–136
Monograph of *Nuphar* (Nymphaeaceae). A, 1–95; keys 24–25, 42, 72–74
Moody, Michael L. 101–107
Morphological characters for grouping *Phragmites* 125–136
Myriophyllum quitense 101–107

Naturalized *Syringa reticulata* 222–224
NEBC Meeting News 111–116, 230–236, 362–365
Nenuphar 1–95; *N. advena* 68, *N. lutea* 25, *N. pumila* 40
New Books 110, 361
New Brunswick 101–107
New England Boneset, *Eupatorium novae-angliae* (Asteraceae). Relationships of the, 145–160
New England flora 237–360, 369–386
New Mexico 395–414
New records: Connecticut 459–463; Maine 430–447; Massachusetts 456–458; Mexico 187–196; New Brunswick 101–107
New reports for Maine: *Fuscopannaria praetermissa*, *Psorula rufonigra*, and *Spilonema revertens* 430–447
New reports for New England: *Buellia ocellata*, *Cladonia symphycarpia* 430–447
New reports for ultramafic soils worldwide, lichens 430–447
New reports of endangered *Chamaedorea* species (Arecaceae) from southeastern Mexico, with notes on conservation status, habitat, and distribution. 187–196
New species 137–144, 395–414, 448–455

New species from the coastal plain of North and South Carolina. ***Eupatorium paludicola*** (Asteraceae): A. 137–144, key 143

New taxa of *Camissonia* (Onagraceae); *Erigeron*, *Hymenoxys*, and *Tetradymia* (Compositae); *Lepidium* and *Physaria* (Cruciferae) from Arizona, New Mexico, and Utah. 395–414

Nomenclature 161–186, 395–414

North America 395–414

North Carolina 137–144

Nummularia 161–186; *N. lanceolata* 172

Nuphar 1–95, taxonomic treatment 22; section *Astylus* 53–86; section *Nuphar* 25–53; *N. advena* 68–82 (key 72–74), ß *tomentosa* 68, subsp. *advena* 74–77 (map 71), subsp. *orbiculata* 78–80 (map 73), subsp. *ozarkana* 77–78 (map 72), subsp. *typica* 70, subsp. *ulvacea* 80–82 (map 73), subsp. *variegata* 54, var. *brevifolia* 69, var. *cubana* 69, var. *hybrida* 82, var. *minor* 82, var. *variegata* 53; *N. affinis* 31; *N. americana* 68, *sensu* Prov. 54; *N. bornetii* 34; *N. centricavata* 34; *N. confusa* 27; *N. intermedia* f. *chlorocephala* 31, f. *luteocephala* 31; *N. japonica* 34–38 (map 36), subvar. *lutea* 34, subvar. *rubrotincta* 34, var. *saijoensis* 38, var. *stenophylla* 34, var. *subintegerrima* 34; *N. jurana* 31; *N. kalmiana* 48; *N. lutea* 25–31 (illus. 26, map 29), ß *genuina* 27, ß *kalmiana* 49, ß *punctata* 27, ß *submersa* 27, f. *parviflora* 27, f. *submersa* 27, f. *terrestris* 28, subsp. *advena* 68, subsp. *orbiculata* 79, subsp. *ozarkana* 77, subsp. *polysepala* 58, subsp. *pumila* 40, subsp. *rubrodisca* 82, subsp. *sagittifolia* 63, subsp. *ulvacea* 80, subsp. *variegata* 54, var. *minor* 27, var. *purpureosignata* 27, var. *rubropetala* 27, var. *urceolata* 27, × *pumila* var. *media* 31, × *pumila* var. *polypetala* 31, × *pumila* var. *sublutea* 31; *N. microphylla* 48–53 (map 50, illus. 51), f. *multisepala* 49; *N. minima* β *spenneriana* 31; *N. minor* 27; *N. oguraensis* 44, var. *akiensis* 45, var. *submersa* 45; *N. orbiculata* 79; *N. ozarkana* 77; *N. ozeensis* 34; *N. polysepala* 58–63 (map 60, illus. 61), var. *picta* 58; *N. pumila* 40–42 (key 42), β *spenneriana* 31, f. *submersa* 34, f. *terrestris* 34, subsp. *oguraensis* 44–47 (map 46), subsp. *pumila* 42–44 (map 45), subsp. *sinensis* 47–48 (map 46), var. *glabrata* 34, var. *hookerii* 34, var. *ozeensis* f. *rubro-ovaria* 34, var. *rehsteineri* 34, var. *timmii* 34, × *lutea* f. *submersa* 32, × *lutea* f. *terrestris* 32; *N. puteora* 70; *N. rivularis* 31; *N. sagittifolia* 63–68 (map 65, illus. 26, 66); *N. schlierensis* 27; *N. sericea* 26, var. *denticulata* 27; *N. shimadai* 34; *N. sinensis* 47; *N. spathulifera* 26; *N. subpumila* 34; *N. systyla* 27; *N. tenella* 26; *N. ulvacea* 80; *N. variegata* 53–58 (map 56); *N.* × *interfluitans* 69; *N.* × *intermedia* 31; *N.* × *rubrodisca* 82–86 (map 84); *N.* × *saijoensis* 38–39 (map 36); *N.* × *spenneriana* 31–34 (map 33)

Nymphaea 1–95; *N. advena* 68, *sensu* Aiton 54, subsp. *erythraea* 69, var. *variegata* 53; *N. arifolia* 68; *N. bombycina* 79; *N. chartacea* 69; *N. fluviatilis* 69; *N. fraterna* 54; *N. hastata* 64; *N. intermedia* var. *hirtella* 31; *N. japonica* 34; *N. kalmiana* 48; *N. longifolia* 64; *N. ludoviciana* 69; *N. lutea* 25, ß *kalmiana* 48, ß *minima* 34, ß *pumila* 40, f. *terrestris* 27, *sensu* L. 34, *sensu* Thunb. 34, var. *harzii* 27, var. *puberula* 27; *N. macrophylla* 68; *N. microcarpa* 69; *N. microphylla* 48–49; *N. orbiculata* 78; *N. ovata* 69; *N. ozarkana* 77; *N. polysepala* 58; *N. puberula* 69; *N. pumila* 40; *N. rubrodisca* 82; *N. sagittaefolia* 63; *N. sagittata* 63; *N.*

sagittifolia 63; *N. ulvacea* 80; *N. umbilicalis* 26; *N. variegata* 53; *N.* × *fletcheri* 82; *N.* × *intermedia* var. *media* 31, var. *polypetala* 31, var. *sublutea* 31
Nymphaeaceae 1–95
Nymphona 1–95; *N. advena* 68–74; *N. lutea* 25–31
Nymphozanthus 1–95; *N. advena* 68; *N. centricavatus* 34; *N. europaeus* 26; *N. japonicus* 34; *N. luteus* 26; *N. microphyllus* 49; *N. ozarkanus* 77; *N. polysepala* 58; *N. rubrodiscus* 82; *N. sagittifolius* 63; *N. variegatus* 54, var. *lutescens* 54; *N. vulgaris* 26
Nyphar 1–95; *N. advena* 68–74; *N. lutea* ß *kalmiana* 48–53

Olday, Fred C. 430–447
Oleaceae 222–224
Onagraceae 395–414
Ontario 125–136
Oswald, W. Wyatt, Barbara C. S. Hansen, and David R. Foster. Comparison of pollen and stomata in late-glacial and early-Holocene lake sediments from eastern Massachusetts. 225–229 (New England Note)
Ozarks 197–221

Pachysandra terminalis, new record for Connecticut 459–463
Padgett, Donald J. A monograph of *Nuphar* (Nymphaeaceae). 1–95
Paleoecology, comparison of stomate and pollen records 225–229
Palms, Mexican 187–196
Parfitt, Bruce D. 222–224
Paris, Catherine A. 369–386
Payne, Richard E. and Bernd Blossey. Presence and abundance of native and introduced *Phragmites australis* (Poaceae) in Falmouth, Massachusetts. 96–100 (New England Note)
Pentagonia 161–186; *P. biflora* 168
Pérez Farrera, Miguel A., Andrew P. Vovides, Carlos Iglesias, Nayely Martínez Meléndez, and Rubén Martínez Camilo. New reports of endangered *Chamaedorea* species (Arecaceae) from southeastern Mexico, with notes on conservation status, habitat, and distribution. 187–196
Persicaria 161–186; *P. opelousana* 176
Phragmites australis (Poaceae) from eastern North America. 415–429
Phragmites australis 96–100, 125–136, 415–429
Phragmites australis subsp. *australis* and subsp. *americanus* 125–136, key 135–136
Phylogeny, *Nuphar* 1–95
Physalis 161–186; *P. carpenteri* 174–175; *P. carpenterii* 174
Physaria 395–414; ***P. arizonica*** var. ***andrusensis***, var. nov. 403–405
Picea glauca and *P. mariana* stomate and pollen records in Massachusetts lake sediments 225–229
Pinus resinosa, Holocene biogeography and genetic diversity 369–386
Plant migration 369–386
Plantago 161–186; *P. caroliniana sensu* Pursh 175; *P. heterophylla* 175; *P. ludoviciana* 175; *P. perpusilla* 175
Poaceae 96–100, 125–136, 415–429
Pollen and stomata in late-glacial and early-Holocene lake sediments from eastern Massachusetts. Comparison of, 225–229 (New England Note)
Polygonella articulata, Holocene biogeography and genetic diversity 369–386
Polygonum 161–186; *P. beyrichianum* 177; *P. chamissonianum* 177; *P. geniculatum* 176–178; *P. meisnerianum* 177–178, var. *beyrichianum* 177–178, var. *jalapense* 177; *P. multiangulare* 177; *P. opelousanum* 176; *P. turkheimii* 177
Polyploidy 145–160, 415–429
Porcher, Richard D. 137–144

Potamogeton 161–186; *P. amplexicaulis* 178; *P. bupleuroides* 178; *P. fulvus* 178–179; *P. loeselii* 178; *P. perfoliatus* 178, var. *bupleuroides* 179, var. *caudiformis* 178, var. *cordaitlanceolatus* 178, var. *densifolius* 178, var. *gracilis* 178, var. *loeselii* 178, var. *mandschuriensis* 178, var. *prolixus* 179, var. *pseudodensus* 178, var. *rotundifolius* 178

Presence and abundance of native and introduced *Phragmites australis* (Poaceae) in Falmouth, Massachusetts. 96–100, map 97 (New England Note)

Prunus sect. *Prunocerasus*, Holocene biogeography and genetic diversity 369–386

Psorula rufonigra, new report for Maine 430–447

Quaternary history of the New England flora 369–386

Quercus 161–186; *Q. andromeda* 179–180; *Q. bumeliaefolia* 180; *Q. carpenteri* 180–181; *Q. carpenterii* 181; *Q. falcata* var. *pagodifolia* 181; *Q. pagoda* 180–181; *Q. pagodifolia* 181; *Q. peckiana* 181; *Q. rhombifolia* 181–182; *Q. rubra* var. *pagodifolia* 181, var. *pogodifolia* 181

Rajakaruna, Nishanta 430–447

Rare North American disjunct, *Myriophyllum quitense* 101–107

Red algal species, ***Tuomeya americana***, comb. nov. 464–467

Refugia 369–386

Refugia and migration in the Quaternary history of the New England flora. 369–386

Relationships of the New England Boneset, *Eupatorium novae-angliae* (Asteraceae). 145–160

Reviewers of Manuscripts 475

Rhynchospora distribution in New England 237–360, maps 345–347

Rhytiglossa 161–186; *R. viridiflora* 169

Riddell, John L., North American taxa published in 1853 161–186

Ropalon 1–95; *R. sagittatum* 63

Sagitta 161–186; *S. graminea* 184

Sagittaria 161–186; *S. lancifolia* var. *papillosa* 183; *S. acutifolia* 184; *S. angustifolia* 184; *S. cycloptera* 184; *S. eatonii* 184; *S. edwardsiana* 184; *S. falcata* 182; *S. gigantea* 182–184; *S. graminea* subsp. *edwardsiana* 184, var. *cycloptera* 184, var. *graminea* 183; *S. lancifolia* subsp. *media* 182, var. *angustifolia* 184, var. *falcata* 182, var. *media* 182; *S. papillosa* 183; *S. paradoxa* 183–184; *S. plantaginifolia* 182; *S. purshii* 184; *S. sagittifolia* var. *simplex* 184; *S. simplex* 184; *S. stolonifera* 184; *S. trachysepala* 183

Saint John River estuary system, New Brunswick 101–107

Salix atrocinerea, new record for Connecticut 459–463

Saltonstall, Kristin, Kelsey Glennon, Alison Burnett, Richard B. Hunter, and Kimberly L. Hunter. Comparison of morphological variation indicative of ploidy level in *Phragmites australis* (Poaceae) from eastern North America. 415–429

Saxifraga oppositifolia, Holocene biogeography and genetic diversity 369–386

Schilling, Edward E. 137–144, 145–160

Schilling, Edward E., Richard J. LeBlond, Bruce A. Sorrie, and Alan S. Weakley. Relationships of the New England Boneset, *Eupatorium novae-angliae* (Asteraceae). 145–160

Schoenoplectus distribution in New England 237–360, maps 348–351

Scirpus distribution in New England 237–360, maps 351–356

Scleria distribution in New England 237–360, maps 356–357

Searcy, Karen B. 456–458
Sedges 237–360
Serpentine ecology 430–447
Sichuan 448–455
Sorrie, Bruce A. 137–144, 145–160
South Carolina 137–144
Specularia 161–186; *S. biflora* 168; *S. californica* 168; *S. ovata* 168; *S. perfoliata* f. *ramosa* 168, f. *rigida* 168
Spilonema revertens, new report for Maine 430–447
Spirillus 161–186; *S. perfoliatus* 179
Springer, Joshua C. and Bruce D. Parfitt. *Syringa reticulata* (Oleaceae) naturalized in northwestern Vermont. 222–224 (New England Note)
Steironema 161–186; *S. ciliatum* var. *angustifolium* 172, var. *heterophyllum* 172; *S. floridum* 171; *S. graminеum* 172; *S. lanceolatum* 172, var. *angustifolium* 172; *S. radicans* 173
Stem color and correlated morphological characters for grouping *Phragmites* (Poaceae) taxa in eastern Ontario. 125–136, key 135–136
Stephenson, Steven L., Harold S. Adams, and Cynthia D. Huebner. Upland forest vegetation of the Ozark Mountains in northwestern Arkansas. 197–221
Stomatal density, *Phragmites* 415–429
Sun, Hang 448–455
Syringa reticulata (Oleaceae) naturalized in northwestern Vermont. 222–224 (New England Note)

Taxonomic assessment of the North American taxa published by John L. Riddell in 1853. 161–186
Taxonomy, *Nuphar* 1–95
Tetradymia 395–414; ***T. canescens*** var. ***thorneae***, var. nov. 405–407
Townsend, J. F. 137–144
Trichophorum distribution in New England 237–360, maps 357–358
Trillium grandiflorum, Holocene biogeography and genetic diversity 369–386
Triodanis 161–186; *T. biflora* 168
Triticum turgidum (*T. dicoccum*), new record for Connecticut 459–463
Tucker, Gordon C. Additions to the flora of Connecticut. 459–463 (New England Note)
Tuomeya 464–467; ***T. americana***, comb. nov. 465; *T. fluviatilis* 465

Ultramafic ecology 430–447
Upland forest vegetation of the Ozark Mountains in northwestern Arkansas. 197–221
Urtica 161–186; *U. alba* 185; *U. aureliana* 184–185; *U. berlandiera* 185; *U. chamaedryoides* 185; *U. gracilescens* 185; *U. gracilis* 185; *U. leucotrys* 185; *U. purpurascens* 185; *U. rubra* 185; *U. verna* 185, var. *gracilis* 185
Utah 395–414
Uxpanapa 187–196

Vaccinium uliginosum, Holocene biogeography and genetic diversity 369–386
Vegetation 197–221
Veracruz 187–196
Vermont 222–224
Vovides, Andrew P. 187–196

Watermilfoil, Andean 101–107
Weakley, Alan S. 137–144, 145–160
Welsh, Stanley L. 395–414
Wetland, *Phragmites* 415–429
Whitson, Maggie K. 161–186
Wikstroemia (Thymelaeaceae) from western Sichuan, China. 448–455
Wikstroemia 448–455; **W. guanxianensis**, sp. nov. 449–451; **W. jiulongensis**, sp. nov. 451–454; key to selected species from Sichuan 454–455

Wilbur, Robert L. and Maggie K. Whitson. Taxonomic assessment of the North American taxa published by John L. Riddell in 1853. 161–186

Wynne, Michael J. Authorship of the freshwater red algal species ***Tuomeya americana*** (Batrachospermaceae). 464–467 (Note)

Yeatesia 161–186; *Y. viridiflora* 169–170

Yeatsia 161–186; *Y. laete-virens* 169

Zhang, Yonghong, Hang Sun, and David E. Boufford. Two new species of *Wikstroemia* (Thymelaeaceae) from western Sichuan, China. 448–455